



Synthesis, characterization and anthelmintic activity of 4-[2'-(5'-nitro) imidazolyl] benzoyl (n-methyl) amino acid derivatives

Ramesh Dhani

Department of Pharmaceutical Chemistry, Oil Technological Research Institute,
Jawaharlal Nehru Technological University, Anantapur, (A.P.) - India

Abstract

Compound containing the imidazole ring are very important in living system, such as vitamin B₁₂ and several pilocarpin alkaloids. Condensation of heterocyclic moieties viz nicotinic acid, thiazole, coumarin, quinoline, furan, imidazole etc. with amino acids and peptides resulted in compounds with potent biological activities. Many of the heterocyclic found to exhibit antifungal, antibacterial, cytotoxic, antineoplastic, insecticidal, anti-inflammatory, tyrosinase inhibitory and melanin production inhibitory activities. Imidazoles have been drawn as promising structural units in the field of medicinal chemistry. Introduction of D-amino acids and N-methylation of amino acids like tyrosine, valine, alanine etc enhanced antimicrobial activity. Hence an attempt is made towards the sythesis of 5-nitroimidazolyl-benzoic acid derivative of N-methyl amino acids and peptide using solution phase technique of peptide synthesis. The synthesized amino acid derivatives studied for anthelmintic activity.

Key-Words: Thiazole, Tyrosine, Quinoline, Imidazole, Valine, Threonine, Alanine


Introduction

A number of synthetic imidazoles such as priscal, privine and antihistamine have physiological properties. The antihistamine azamycine is a 2-nitroimidazole. Amongst synthetic imidazoles in use as therapeutic agents are cimetidine for the treatment of peptic ulcer, metronidazole an antibacterial and an antiprotozoal used in the treatment of amoebic dysentery. Miconazole, clotrimazole, ketaconazole are some currently used antifungal imidazole derivatives. They act by inhibiting ergosterol synthesis which is a primary cellular sterol of fungi cell wall. Ketaconazole was the first successful orally active imidazole.

The method includes the introduction of tert-butyloxy carboxyl group (Boc) to amino acids to protect the amino group forming Boc- amino acids. The protection of carboxyl group was done by converting the amino acids into corresponding methyl ester. N-methylation was done by treating with methyl iodide and sodium hydride.

The ester group was then removed by lithium hydroxide. The Boc(N-methyl) amino acids or Boc(N-methyl)dipeptide were coupled to amino acids or Boc(N-methyl)dipeptide were coupled to 4-(2'-(5'-nitro imindazolyl] benzoic acids. The synthesized compounds will be tested for their biological activitie as anthelmintic activitie[1,2].


Some noteworthy works done are mentioned below:
Srivastava R. et al., Worked on 2,5-disubstituted benzimidazole pyrido[1,2-a]benzimidazoles and some tetrasubstituted benzophenones as probenzimidazoles. This compound showed anti- filarial activity against *L.carinii* and *B.malayi*[5].




* Corresponding Author

E.mail: dhaniramesh1@gmail.com

Usha Rani T. et al., Worked on the synthesis of the substituted-9- hydroxy-10-undecyl [2',3',4,5]-thiazol-[2,3-b] benzimidazole-8,11-diones. These compounds where found to possess fungicidal activity against *F.oxysporum*, *Lanata* and bactericidal activity *B.substilis* *E.coli* and *P.vulgaris*[6]



Renukadevi Patil et.al., Worked on the synthesis of 1,2-di-substituted-4-[(5'-substituted-2'-phenylindol-3'-yl)methylene]iminolin-5-(4H)-ones. The compounds were found to exhibit antibacterial activity[7].



Material and Methods

General Method of Amino acid Synthesis:

The method involved in the synthesis of 4-[2'-(5'-nitro)imidazolyl]benzoyl(N-Me) amino acids is the coupling reaction between amino acid and 4-[2'-(5'-nitro)imidazolyl]benzoic acid

Synthesis of N-methyl amino acids methyl ester:


Amino acids were converted into the corresponding methyl ester hydrochloride using thionyl chloride and methanol. The amino end was then protected by introducing Boc-group using di-tert-butylpyrocarbonate and triethylamine to get Boc-L-amino methyl ester. N-methylation of this compound was done by treating with methyl iodide and sodium hydride (Benion method) to get Boc-(N-Me) amino methyl ester [1, 5].

Preparation of 4-[2'-(5'-nitro)imidazolyl] benzoic acid :

A mixture of p-amino benzoic acid (34.25 gms, 250 mmol), dilute hydrochloric acid (15%, 120ml) and water (150ml) was heated to get a clear solution. The solution was cooled to RT and diazotized by the addition of sodium nitrite solution (30%, 48ml). The diazonium salt solution was filtered and to the filtrate, dilute HCl (100ml) and nitroimidazole (250 mmol) and aqueous cupric chloride (5gms in 20ml of water) were added with stirring. Stirring was continued for 6 hrs and kept overnight in the refrigerator. The separated solid was collected by filtration and washed with water. The crude compound was crystallized from acetone to obtain pure of 4-[2'-(5'-


nitro)imidazolyl]benzoic acid[3,6].

SCHEME-1



Preparation of 4-[2'-(5'-nitro)imidazolyl]benzoyl(N-Me)-amino acid

To the (N-Me) amino acid methyl ester (7.0 mmol), THF (20ml), added 4-[2'-(5'-nitro)imidazolyl] benzoic acid (1.631gms, 7.0 mmol), DIPC, Et3N (2.8ml) and stirred at room temp. for 24hr. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure, residue was dissolved in CHCl3, washed with 10% NaHCO3 (10ml) and 5% HCl (10ml), dried over anhydrous Na2SO4 and evaporated under vacuum to get the title compounds. The crude product was recrystallized from CHCl3 and n-hexane [8].



R = Valine, Phenyl alanine, Tyrosine, Threonine, Leucine

Preliminary Analysis of the sample

Thin-layer chromatography (TLC) was commonly used in the qualitative description of the complexity and composition of chemical mixtures.

Application of sample on TLC plates

The sample was applied to the chromatogram by repeated "spotting" above 1-2 cm from one end of the plate with a capillary tube.

The most important precaution was not to apply spots below the level of the top of the solvent system in developing chamber

Developing solvent systems

Development chamber was used for developing chromatogram. Chloroform: Methanol: Water 5:3:2 was the solvent system used for running TLC of these compounds.

Visualization of chromatogram

After developing, the TLC plates were dried and then exposed to iodine vapours in a chamber, since chromatograms of many synthetic products were frequently observed by iodine vapors. R_f value was noted down. Purity of all the synthesized compounds including intermediates was checked by TLC on silica gel G plates. All compounds have shown only single spot indicating the completion of the reaction and the purity of the product obtained.

Evaluation of Anthelmintic activity

Anthelmintics are therapeutic agents used to destroy parasitic worms or remove them from the infected host. The ultimate test of anthelmintic activity is the ability of a chemical agent to eliminate the worms from a specifically parasitized animal with a minimum of toxic effect to the host. A suitable in vitro test can be considered as a useful screening method, although in vivo screening methods provide a natural environment for the studies.

General Procedure:

Anthelmintic activity studies were carried out against earthworms (*Eudrilus eugeniae*) by Garg's method.⁴¹ Suspensions of the samples were prepared by triturating the samples with 15% tween 80 and distilled water and the resultant mixtures were stirred using a mechanical stirrer for 30 mins. The resulting suspensions were used for the activity studies. The suspensions were diluted to contain 100 mg in 5 ml of the test samples. Standard drug, Mebendazole was also prepared with the same concentration in a similar way. Five earthworms of similar sizes were placed in a petri plate of 4 inches diameter containing 50 ml of suspension the test standard drugs (Mebendazole) at RT. Another set of five earth worms was kept as control in 50ml suspension of distilled water and 15% tween 80. 50 ml each of the suspensions of the test compounds were added into separate petri plates containing five earthworms in each. The time required for the paralysis and death of the worms was noted. The death time was ascertained by placing the

earthworms in warm water at 50°C, which stimulated the movement if the worm was alive. Physical data of this activity is shown in table no.7.

Results and Discussion

The new synthesized amino derivatives were screened for anthelmintic activity; the results of anthelmintic activity against *Eudrilus eugeniae* are shown in Table 7

Conclusion

The new 4-[2'-(5'-nitro) imidazolyl] benzoyl (N-Me) amino acid derivatives was synthesized and characterized by physicochemical analytical data. By this studies find the structure-activity relationship and to optimize the structure. The synthesized amino acid derivative i.e., 4-[2'-(5'-Nitro) imidazolyl] benzoyl(N-Me) amino acid derivatives was confirmed by physicochemical & spectral analysis and further screened for anthelmintic activity, all derivatives shows that good activity.

Acknowledgement

In the first place we would like to record my gratitude to my parents and my sister for their inspiration and encouragement given to me during this work with deep appreciation for their determination and enthusiasm at each and every front of my life to transform my dreams into reality. I am very thankful and prevail age to my deep sense of gratitude to Abdul Mohammed Bari, M.pharm, Ph.D, Director of Bright laboratories, Hyderabad.

References

1. Bansal RK, 'Synthesis, Reaction and Mechanism of Heterocyclic Chemistry', *Indian Journal Chemistry*, vol.12, Pg no 34-36, 1996.
2. Belagali SL, Harish K, Boja P and Himaja M., "Synthesis and biological studies of 5 (p-chlorophenyl)-furan-2-carbonyl peptides and 4-[2'-(5'-formyl) furyl]benzoyl peptides", *Indian Journal Chemistry*, vol.6, pg no.378-387, 1998.
3. Belagali SL and Himaja M, "Synthesis and biological activity of pyridine-3- carboxyl amino acids and peptides", *Indian Journal Heterocycle Chemistry*, vol.22, pg no.8-11, 1998.
4. Stanchev M, Tabakova S, Vedenov VG, Galovinsky E, Jung G, "Synthesis and oxazole moieties", *Arch Pharm Weihen*, vol.19, pg no.332 – 292, 1999.
5. Boja P, Belagali SL, Harish K, Holla BS and Gonsalves R. "Synthesis and characterization of some amino acids derivatives of 3-carbomethoxy-methyl-7- hydroxy-4 methyl coumarins", *Indian Journal Heterocycle Chemistry*, vol.3,

- pg no.263, 2000.
6. Belagali SL, Mathew T, Himaja M, Kocienski P. "A highly efficient method of N-Methylation for the amino acid derivatives" *Indian Journal Chemistry*, vol.3, pg. no 45, 1995.
 7. Benoiton NL, Akyusekli D, Chen. FM, "N,N-dialkoxycarbonyl amino acid from the sodium hydroxide mediated reaction of alkyl chloroformates with anhydrides of N-alkoxy carbonyl amino acids", *International Journal of Protein Research*, vol 45, pg no 466-470, 1995.
 8. Bauer AW, Kirby WM, Shersis JC, Turck M, "Antibiotic susceptibility testing by a standardized single disk method" *Indian Journal of chemistry*, vol 45, pg no 493-496, 1996.
 9. Himaja M. Rajiv T, Geetha P, Harish K, Boja et al." hymenamide A, Unusual Biologically active Cyclic Peptide". *Boll Chim. Farmaceutico-Anno*, vol.13, pg.no168-175, 1999.
 10. Ahluwalia VK, Mallika N, Singh RP, Khandhur C, "A convenient synthesis of 3-(4-arylthiazole-2-ylthio)coumarins", *Indian Journal Chemistry*, vol.29, pg no. 667, 1990.
 11. Cosar and Brown, "Antiparasitic drugs IV 2-(4-thiazolyl)benzimidazole", *Indian Journal Heterocycle Chemistry* vol , vol 15, pg-no.1764-1765, 1990.

Physicochemical analysis

Table 1

S/No.	4-[2'-(5'-nitro)imidazolyl] benzoyl(N-Me)Tyrosine	
1.	Mol. Formula	C ₂₀ O ₆ N ₄ H ₁₈
2.	Mol. Weight	410
3.	Melting Point	295°C
4.	Physical state	Brown Solid
5.	R _f Value	0.55
6.	Solvent System	CHCl ₃ : CH ₃ OH : H ₂ O (5:3:2)

Table 2

S/No.	4-[2'-(5'-nitro)imidazolyl]benzoyl(N-Me)Phenyl alanine	
1.	Mol. Formula	C ₂₀ O ₅ N ₄ H ₁₈
2.	Mol. Weight	394
3.	Melting Point	173°C
4.	Physical state	Sticky Brown Mass
5.	R _f Value	0.64
6.	Solvent System	CHCl ₃ : CH ₃ OH : H ₂ O (5:3:2)

Table 3

S/No.	4-[2'-(5'-nitro)imidazolyl] benzoyl(N-Me)threonine	
1.	Mol. Formula	C ₁₅ O ₆ N ₄ H ₁₈
2.	Mol. Weight	350
3.	Melting Point	173°C
4.	Physical state	Brown Solid
5.	R _f Value	0.57
6.	Solvent System	CHCl ₃ : CH ₃ OH : H ₂ O (5:3:2)

Table 4: Physical data of 4-[2'-(5' nitro) imidazolyl] benzoic acid

S/No.	Product Name	Physical state	M.P(⁰ C)	%yield
1.	4-[2'-(5'-nitro)imidazolyl]benzoic acid	Pale brown solid	256	21.49

Table 5: Physicochemical analysis of 4-[2'-(5'-nitro) imidazolyl] benzoyl(N-Me)valine

S/No.	Physicochemical analysis of 4-[2'-(5'-nitro)imidazolyl] benzoyl(N-Me)valine	
1.	Mol. Formula	C ₁₆ O ₅ N ₄ H ₁₈
2.	Mol. Weight	346
3.	Melting Point	120°C
4.	Physical state	Reddish Brown solid
5.	R _f Value	0.53
6.	Solvent System	CHCl ₃ : CH ₃ OH : H ₂ O (5:3:2)

Table 6

S/No.	Physicochemical analysis of 4-[2'-(5'-nitro)imidazolyl] benzoyl(N-Me)leuline	
1.	Mol. Formula	C ₁₇ O ₅ N ₄ H ₂₀
2.	Mol. Weight	360
3.	Melting Point	120°C
4.	Physical state	Light Brown Solid
5.	R _f Value	0.55
6.	Solvent System	CHCl ₃ : CH ₃ OH : H ₂ O (5:3:2)

Table 7: Data of anthelmintic activity

S/No.	Compound	Concentrations (mg)	Mean paralyzing time (min) + S.E	Mean death time (min) +S.E
1.	Control	-	N.E	
2.	Mebendazole	100	6.00 ± 0.34	7.40
3.	1	100	6.90 ± 0.31	7.35
4.	2	100	6.12 ± 0.35	7.34
5.	3	100	6.10 ± 0.51	6.90
6.	4	100	7.04 ± 0.32	8.29
7.	5	100	6.10 ± 0.31	7.40

S.E represents Standard Error and N.E. indicates No Effect