



## Screening of *Camellia oleifera* Abel. seeds extract for free radical scavenging activity

Ranjit Singh\* and C. Sankar

KMCH College of Pharmacy, Department of Pharmaceutics, Coimbatore, (TN) - India

### Abstract

The main objective of the study was to find the antioxidant value of *Camellia oleifera* Abel seeds. Antioxidants have been reported to prevent oxidative damage caused by free radical and can be used in cardiovascular and anti-inflammatory diseases. The free radical scavenging activity of ethanolic extract of *Camellia oleifera* Abel seeds was studied with the help of a stable DPPH radical. Anti-oxidant activity was measured by 1, 1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging assays method and total phenolic and tannin content was also determined. The ethanolic extract was found to have strong free radical scavenging capacities, with IC<sub>50</sub> values lower than 0.8 mg/ml.

Key-Words: *Camellia oleifera* Abel; Seeds; Free radical scavenging activity; Antioxidant, Phenols; Tannins

### Introduction

The DPPH free radical scavenging assay is simple and widely used for bioactive compound discovery. In conjunction with a microtitre plate reader, the assay can be easily carried out as a high throughput test<sup>1</sup>.

### Material and Methods

#### Chemicals

1, 1-Diphenyl-2-picrylhydrazyl (DPPH) was bought from Aldrich (Shanghai). Analytical grade methanol was from Ranbaxy chemicals Pvt. Ltd (Mumbai). A Multiscan (Ascent) microtitre plate reader was purchased from Thermo Bio analysis Company (Finland).

#### Collection of Plant materials

The seeds of the plant belonging to the genus *Camellia* (Theaceae) were collected from Korakundah, Nilgiri estate (Tamil Nadu) and were authenticated. The seeds were used for the extraction and anti scavenging activity.

#### Method of extraction<sup>2</sup>

The seeds were dried and made into fine powder by grinding and extracted for 48 hrs at room temperature with ethanol. For every milligram of seeds powder, 10 ml of ethanol were added. Extraction was carried out under shaking with 80 rev/min. The suspensions were centrifuged, and the supernatant diluted at different concentrations, which were used as test samples.

### Quantitative analysis

Quantitative measurement of radical scavenging properties was carried out in a 96 wells microtitre plate assay<sup>1</sup>. The reaction mixture contained 100 µl of test samples (or ethanol as a blank) and 100 µl of a 1 mM solution of DPPH in EtOH. 100 µl of 10 mM solutions of ascorbic acid was used as positive control. Decoloration was measured at 517 nm after incubation for 20 min. Measurements were performed at least in triplicate. The actual decrease in absorption induced by the test compounds was compared to that of the positive controls. IC<sub>50</sub> values calculated denote the concentration of sample required to scavenge 50% of DPPH radical.

### Results and Discussion

The ethanolic extract of *Camellia oleifera* Abel seeds were screened for DPPH radical scavenging activity<sup>1</sup>. The extract had shown strong free radical scavenging capacities, with IC<sub>50</sub> values lower than 0.8 mg. The activity of the extract is due to the active constituent present in the seeds, Theanine. The seeds contain tannins and other polyphenols such as flavonoids. Theanine is responsible for the strong free radical scavenging activities<sup>3</sup>. It is well known that tannins have anti-inflammatory, antidiarrheic and wound healing properties. Phenols such as methyl gallate possess antimicrobial properties, and physiological redox processes in the microbial cells may be disrupted by the strong reducing activity of tannins<sup>4</sup>. The anti-inflammatory properties of tannins are mainly due to their free radical scavenging activities, as free radical

### \* Corresponding Author

E.mail: ranjitsingh08685@gmail.com  
Mob.: +91-9042978206

scavengers can stop some of the processes of inflammatory response<sup>5</sup>.

### **Conclusion**

The ethanolic extract of *Camellia oleifera* Abel seeds was found to have strong free radical scavenging capacities, with IC<sub>50</sub> values lower than 0.8 mg/ml.

### **Acknowledgement**

The authors are thankful to The Principal Dr. A. Rajasekaran, KMCH College of Pharmacy for providing the necessary facilities in the College. Sincerely thanks to Dr. C. Sankar, professor Department of Pharmacuetics, KMCH College of Pharmacy, Coimbatore -35, Tamil Nadu (INDIA) for his valuable support. We would also like to thank our colleagues, lab assistants for their support.

### **References**

1. Cotellet JA, Bernier JL, Catteau JB, Pommery J, Wallet JC, Gaydou EM. Free Radical Biol Med 1996; 20:35.
2. *Journal of Medicinal Plants Research*, volume 6, number 13, 9 April 2012; ISSN 1996-0875.
3. Huang TK, Ding ZZ, Zhao SX. Modern compendium materia medica. Beijing: Chinese Medical Science and Technology Press, 2001. p. 1.
4. Lin HC, Ding HY, Wu YC. J Natl Prod 1998; 61:343.
5. Sun CP, Zhang JZ, Duan SJ. Free radical biology. Anhui: Chinese Science and Technology University Press, 1999.
6. Hirano R, Sasamoto W, Matsumoto A. J Nutr Sci Vitaminol 2001; 47:357.
7. Yokozawa T, Chen CP, Dong E. Biochem Pharmacol 1998; 56:213.
8. Yao XS. Natural products chemistry. Beijing: Chinese Health Press, 1997. p. 241.
9. Chen HF. Dictionary of plants bioactive compounds. Beijing: Chinese Medical Science and Technology Press, 2001.
10. Bruneton J. Pharmacognosy, phytochemistry and medicinal plants. Paris: Lavoisier Publishing, 1995. p. 1.
11. Liu F, Ng TB. Life Sci 2000; 66:725.
12. Mi TY, Li CQ, Zhou HR, Qian XH. Anhui economical plants flora Anhui. Anhui Science and Technology Press, 1999.
13. Salatino A, Salatino ML, Giannasin DE. Biochem Syst Ecol 2000; 28:545.
14. Takeda Y, Okada Y, Masuda T, Hirata E, Shinzato T, Takushi A, Yu Q, Otsuka H. Chem Pharm Bull 2000;48:752.
15. Piacente S, Pizza C, De Tommasi N, Mahmood N. J Natl Prod 1996;59:565.
16. Miliauskas, G., P.R. Venskutonis and T.A. Van Beek, 2004. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. Food Chem., 85: 231-237.

**Fig. 1. *Camellia oleifera* Abel plant.**



**Fig. 2. *Camellia oleifera* Abel seeds**

