



INTERNATIONAL JOURNAL OF PHARMACY & LIFE SCIENCES

1, 5, 8-Trihydroxy-3-methoxyxanthone from *Swertia corymbosa* (Griseb.) Wight ex.C.B.Clarke

A.Saraswathy and S.Ariyanathan*

Captain Srinivasa Murti Drug Research Institute for Ayurveda (CCRAS)
Arumbakkam, Chennai, (TN) - India

*Research Scholar, SASTRA University, Thanjavur, (TN) - India

Abstract1,5,8-Trihydroxy-3-methoxyxanthone was isolated from *Swertia corymbosa* (Griseb.) Wight ex.C.B.Clarke**Key-Words:** *Swertia corymbosa*, 1,5,8-Trihydroxy-3-methoxyxanthone**Introduction**

Swertia corymbosa (Griseb.) Wight ex.C.B.Clarke (Gentianaceae) is known as Poovainagai in Tamil¹ is used in the traditional systems of medicine as an antidote for poisons, diarrhoea, fever and as a stomach wash in cattle². Friedelin, epi-friedelinol, mangiferin, α -amyrin, lupeol, oleanolic acid, ursolic acid³ were reported from the n-hexane and chloroform extracts of the plant. Further work on the methanol extract yielded yet another xanthone 1, 5, 8-trihydroxy-3-methoxyxanthone. This is the first report of the compound from this plant.

Material and Methods**Collection of plant material**

The plant material was collected from Thirunelveli District, Tamil Nadu in the month of January and identified by Dr M.B. Vishwanathan Dept. of botany, of Sri Paramakalyani centre for environmental sciences, Manonmaniam Sundaranar University of Tamilnadu. A voucher specimen (MBV and NR 6759) has been deposited in the herbarium of the centre.

Extraction and isolation

The shade dried plant material (1kg) was coarsely powdered and extracted successively with n-hexane, chloroform and methanol. The extracts were concentrated under vacuum after distilling off the solvent on a water bath. The methanol extract (60g) was suspended in water (1000 ml) and partitioned successively in a separating funnel with diethyl ether (250ml), ethyl acetate (250ml) and n-butanol(100ml). Ethyl acetate soluble fraction after drying over anhydrous sodium sulphate and concentration (1.5g) was column chromatographed over silica gel (acme, 100-200 mesh, 1:22). Elution of the column with chloroform - ethyl acetate (9:1) afforded a solid which was crystallised from chloroform to get the pure compound as pale yellow amorphous powder. M.p 212 -214°C. UV λ max (MeOH) nm 210, 239, 268, 328. IR ν_{max} (KBr) hydroxyl (3459 cm^{-1}), chelated carbonyl (1661, 1644 cm^{-1}) and aromatic system at 1608, 1581, 1496, 838-650 cm^{-1} . ^1H NMR $\delta(\text{CDCl}_3)$: 11.96, 11.87 (1H each, br s, 1-OH and 8-OH), 7.30 (1H, d, J=9.0Hz, 7-H), 6.84 (1H, d, J=9.0Hz, 6-H), 6.34 (1H, d, J=2.3Hz, 2-H), 6.41 (1H, d, J=2.3Hz, 4-H), 5.38 (1H, s, 5-OH), 3.89 (3H, s, 3-OCH₃).


Results and Conclusion

The pale yellow compound isolated from the ethyl acetate fraction appeared brown on exposure to iodine vapour, UV 366nm and with vanillin sulphuric acid (m.p. 212 - 214°) was analyzed for C₁₄H₁₀O₆ (M⁺ 274). It gave a green colour iron (III) chloride. The UV spectrum of the compound run in MeOH showed λ max 210, 239, 268, 328 nm, suggesting it to be a xanthone. Addition of AlCl₃ shifted the bands at 268 and 328 nm to 287 and 362. However, addition of con. HCl did not affect the shifts much. This indicated peri interaction of

*** Corresponding Author:**

E-mail: ariy78@gmail.com

phenolic OH with carbonyl. Neither bathochromic nor hypsochromic shifts were observed in the UV spectrum recorded in the methanolic sodium acetate indicating the absence of free hydroxyl at C-3. Its IR spectrum had bands for the presence of xanthone hydroxyl (3459 cm^{-1}), chelated carbonyl ($1661, 1644\text{ cm}^{-1}$) and aromatic system at $1608, 1581, 1496, 838\text{--}650\text{ cm}^{-1}$. In the $^1\text{H NMR}$ spectrum two aromatic protons with meta coupling ($J=2.3\text{ Hz}$) appeared at δ 6.34 (C2 - H) and 6.41 (C4 - H). Two ortho coupled aromatic proton appeared at doublet 7.30, 6.84 with $J=9.0\text{ Hz}$ corresponding to C7- H, C6- H. The methoxy methyl protons appear as singlet at 3.89. The broad singlet at 5.38 indicated the presence of C5- hydroxyl. Two chelated hydroxyls (C1-OH and C8-OH) appeared at 11.96 and 11.87 as broad singlet. The physical and spectroscopic data were comparable with those reported for 1, 5, 8-trihydroxy, 3-methoxy xanthone⁴⁻⁹. From the above evidences the compound was identified as 1, 5, 8-trihydroxy, 3-methoxy xanthone.



Acknowledgements

Thanks are due to The Director General, Central Council for Research in Ayurveda and Siddha, New Delhi for financial support and Prof. Joseph D. Connolly for providing the spectral data. Dr M.B. Vishwanathan, Botanist of Sri paramakalyani centre for environmental sciences, Manonmaniam Sundaranar University of Tamilnadu is gratefully acknowledged for the plant identification.

References

1. Kothawade S. N. Kadam N. R. Aragade P. D. Clarke C.B. (1883). Gentianaceae. In: Hooker J. D. (Ed.) *Flora of British India*, Vol. IV. London: L. Reeve & Co, 126.
2. Abraham Z. (1981). Ethnobotany of the Todas and the Kotas and the Irulas of Nilgiris. In: Jain S.K. (Ed.) *Glimpsee of Indian Ethnobotany*. New Delhi: Oxford & IBH publishing Co.
3. Ramesh N., Vishwanathan M.B., Saraswathy A., Balakrishna K., Brindha P. and Lakshmanaperumalsamy P. (2002). Antimicrobial and phytochemical studies of *Swertia corymbosa*. *Fitoterapia*, **73**:160-164.
4. Purev O., Kh. Oyun, Odontuya G., Tankhaeva A. M., Nikolaeva G. G., Khalid M. Khan, Syed Tasadaque Ali Shah and Wolfgang Voelter. (2007). Isolation and structure elucidation of two new xanthenes from *Gentiana azurium* Bunge (Fam. Gentianaceae).
5. Odontuya G., Kojima K., Rodriguez S., Purev O., Ogihara Y. and Hostettmann K., (1998). *Chem. Pharm. Bull.*, **46**:1827-1828.
6. Khadga S, Khetwal, Binita and Bisht R.S. (1990). Tri and Tetraoxygenated Xanthenes from *Swertia petiolata*, *Phytochemistry*, **29**(4): 1265-1267.
7. Jie Shen and Jun-Shan Yang (2006). Two new Xanthenes from the stem of *Garcinia cowa*. *Chem. Pharm. Bull.*, **54**(1):126-128.
8. Mehtab Parveen and Nizam Ud-din Khan (1988). Two Xanthone glycosides from *Gentiana lutea*, *Phytochemistry*, **27**(11): 3696-3699.
9. Takaaki Hayashi and Takashi Yamagishi (1998). Two Xanthenes glycosides from *Gentiana lutea*. *Phytochemistry*, **27**(11): 3696 - 3699.