



**Screening of potential efficacy of dietary ginger on ethanol induced oxidative stress in rat cardiac tissue: A study on changes in basic metabolic profiles**

G.Venkata Subbaiah\*, R. Naliniprasad, J. Himabindu, M.Anila Devi,  
I. Nagaraju, P. Chandrakala, P. Mohan., K. R. Shanmugam, CH. Ramakrishna, S. Ravi, B.  
Shanmugam and K. Sathyavelu Reddy\*

Division of Molecular Biology and Exercise Physiology, Department of Zoology Sri Venkateswara University, Tirupati, (AP) - India

**Abstract**

The present study was premeditated to examine the possible mechanisms where by ginger (*Zingiber officinale*) could protect cardiac tissue from alcohol toxicity in rats. The carbohydrate metabolic profiles like total carbohydrates, pyruvate, total proteins, free amino acids and lactate levels were measured in heart tissue. The total carbohydrates, pyruvate, and total proteins were significant declined while free amino acids, lactate levels were significant increased in alcohol intoxicated rats. Whereas with ginger (200 mg/kg body weight) treatment shown significant increase in the total carbohydrates, total proteins and pyruvate levels, whereas free amino acids, lactate levels were significant drop in the cardiac tissues. From the present study, we conclude that ginger protects the heart tissue from alcohol toxicity in rats, this may be due to the presence of many bioactive compounds in ginger.

Key-Words: Alcohol, Ginger, Heart, Carbohydrate metabolic profiles, Rats

**Introduction**

Worldwide alcohol has become the most socially accepted addictive drug (Guo, & Ren, 2010). Although, alcohol dependence is a serious medical illness (American Psychiatric Association, 1994). Alcohol constitutes a substantial health and economic burden, costing an estimated \$184 billion in expenditures stemming from alcohol-related chronic diseases such as heart disease (George, & Figueredo, 2010), liver disease (Cederbaum, *et al.*, 2009), cancer (Seitz, & Becker, 2007), diabetes (Baliunas, 2009). Alcohol is absorbed by the gastrointestinal tract at various levels (Holford, 1987). Moderate alcohol intake has been shown significant protective effect against vascular disease (Mukamal *et al.*, 2005). These effects can probably be attributed to the potential for alcohol to increase protective HDL cholesterol levels and decrease platelet aggregation. In contrast, alcohol is generally accepted to be a toxic compound when it take excessive results its metabolized by alcohol dehydrogenase to produced disproportionate amount of acetaldehyde by the microsomal ethanol oxidizing system in the liver (Ohkubo *et al.*, 2009).

**\* Corresponding Author**

E.mail: ganjikuntasubbaiah@yahoo.com,  
ksreddy2008@hotmail.com

Lieber (1995) reported that acetaldehyde; one of the products in alcohol metabolism alters the redox state in the cytosol, which in turn responsible for the abnormalities in antioxidant, lipid metabolism and carbohydrate metabolism. There is evidence that reactive oxygen species which are produced in alcohol toxicity (Robin, *et al.*, 2005) alters the mitochondrial structural, function in various organs including heart (Regan, 1990), liver (Klein, & Harmjanz, 1975). Since ancient times, medicinal plants used to cure various ailments in humans (Butt *et al.*, 2009). A number of scientific investigations have led to the recognition of a safe status for such natural products. In Chinese, Ayurveda, Unani-Tibb like traditional medicines ginger have been used as medicinal plant (Rong *et al.*, 2009). Ginger posses various pharmacological activities including hypoglycemia, anticancer, anticardiac antirenal and hepatoprotective and antioxidant (Nicoll and Henein, 2009). Ginger has many antioxidant compounds; these compounds may either mitigate or prevent generation of free radicals in toxic conditions. The active ingredients of ginger include gingerols, shagogals, phytochemicals and other compounds show antioxidant activity in various models (Dugasani *et al.*, 2010). The present study was

design to explore the cardiac protective property of ginger ethonolic extract against alcohol induced cardiac toxicity in the rats.

### Material and Methods

#### Ginger extracts preparation

The rhizomes of *Zingiber officinale* were purchased from local market in Tirupati with authenticated by botanist in the department of Botany, S.V. University, and Tirupati. The rhizomes were shade dried at room temperature and were crushed to powder. 120g of powder has taken and macerate in 1000 ml of 95% ethanol for 12 h at room temperature, then filtered and squeezed with muslin cloth to obtain ethanol extract juice. This process was repeated three times and finally collection of this juice were dried in rotary evaporator (Model:HS-2005V) from this we had get jelly and then this jelly was converted to powder in lyodel freezer. We has done dose dependent studies by using, 50 mg/kg, 100 mg/kg, 150 mg/kg, 200 mg/kg, 250 mg/kg and 300 mg/kg. of this 200 mg/kg dose showed good antioxidant activity. So this study we selected dose of 200 mg/kg of ethanol extract of ginger.

#### Animals and treatments

The study was approved by Institutional Animal Ethics Committee (IAEC) [Reg No: 10/i/a/CPCSEA/IAEC/SVU/KSR-GVS/dt 15/11/2011]. Twenty four Wistar male albino rats weighting  $200 \pm 25$  g were used. They obtained from Indian Institute of Science, Bangalore. Animals were kept in laboratory at constant room temperature ( $26 \pm 2^{\circ}\text{C}$ ) at least one week before and throughout the experimental period. Commerical pellet diet and water were provided *ad libitum*. After acclimatization 24 rats were divided into four groups of 6 animals each and treated as follows: **Group I: Normal control (NC):** rats received 2 % of Tween- 80 in normal saline. **Group II: Ginger treatment (Gt) :** rats received ethanol extract of ginger (200 mg/kg body wt) orally for 30 days. **Group III: Alcohol treatment (At) :** rats received alcohol at dose of 2g/kg for 30 days. **Group IV: Ginger+Alcohol treatment (At+Gt) :** rats received ginger for 30 days followed by alcohol (2g/kg) for 30 days. At the end of 30 days treatment period, the rats were sacrificed by cervical dislocation and heart tissues were isolated, washed with ice cold saline, immersed in liquid nitrogen and stored in deep freezer at  $-80^{\circ}\text{C}$  for further biochemical analysis. The carbohydrate metabolic profiles such as total proteins, total carbohydrates, total free amino acids, pyruvate and lactate levels were estimated in cardiac tissue by the methods of Lowry *et al.*, 1951, Carroll *et al.*, 1956, Moore and Stein, 1954, Friedmann and Hangen, 1942 and Barker and

Summerson (1941) as modified by Huckabee 1961 respectively.

#### Chemicals

In the present study all chemicals used were of Analar Grade (AR) and purchased from the following scientific companies: Sigma (St.Louis, MO, USA), Fischer (Pitsburg, PA, USA), Merck (Mumbai, India), Ranbaxy (New Delhi, India).

#### Statistical analysis

The data has been analyzed by using SPSS (Version 16.0; SPSS Inc., Chicago, IL, USA) and M.S. Office, Excel Software for the significance of the main effects (factors), and treatments along with their interactions. The data has been compared using one way ANOVA with Dunnett's multiple comparison test and differences were considered significant at  $p < 0.05$ .

#### Results and Discussion

Ginger phenol compounds like gingerols, shagogs Possesses antioxidant properties due to their free radical scavenging activities. (Stadtman, 2004; Siddaraju & Dharmesh, 2007; Shanmugam *et al.*, 2010). Ginger has been used to cure diarrhea, dysentery, fever, cough, ulcers boils and wounds (Young, HV *et al.*, 2005). Ajith *et al.*, (2007) reported that ginger ameliorat cisplatin- induced nephrotoxicity. A combination of ginger and garlic has been reported to produce hypoglycemic and hypolipidemic effects. In another study ginger has protected the tissue from oxidative stress.

The data obtained from our study demonstrate that total proteins (Figure D) were significantly decreased in the alcohol treated rats than normal control group; meanwhile total proteins were increased in the ginger treated group compared to ethanol group. Proteins are most important to maintain the structure of the body. Feinman, 1998 & Lieber, 2003 reported that in alcohol subjects, protein levels are decreased. Chronic exposure of rats to a diet containing ethanol inhibit the protein synthesis as result of declined messenger RNA translation consecutively 25% loss in cardiac protein per heart (Vary *et al.*, 2001). The major ethanol metabolite acetaldehyde may contribute to cardiac dysfunction, hypertrophy and heart failure by either its direct toxicity or promoting elevated levels of catecholamines and reactive oxygen species (ROS) (Zhang *et al.*, 2010). Although, alcohol metabolite acetaldehyde has cytotoxic effect within the cells or tissues due to reacting with nucleic acids, proteins, peptides, amino acids, lipids and carbohydrates (Bartsch and Nair, 2000). Other early reports also implicated enhance lipid peroxidation or generation ROS could results reduces the tissue protein synthesis (Matias *et al.*, 1999). On the other hand total protein



levels are increased in ginger treated and ginger treatment in alcoholic rats. Many reports confirmed that ginger's antioxidant activity. Siddaraju and Dharmesh (2007) reported that ginger-free phenolic and ginger hydrolysed phenolic fractions exhibited free radical scavenging activity. Thus ginger prevented the oxidative modification of proteins by its antioxidant property.

The variations in carbohydrates and pyruvate levels in cardiac tissue are shown in figure B and C. There was a significant decrease ( $P < 0.05$ ) in the carbohydrate and pyruvate levels in ethanol treated rats when compared to normal control rats. Alcohol metabolites acetaldehyde, acetate and their end product acetyl CoA levels were increases NADH/NAD<sup>+</sup> ratio, which reflects to deplete the glycogen levels due to a blockage of gluconeogenesis in the hepatic tissue (Lieber, 1991). Randle *et al.*, (1988) has confirmed that acetyl CoA inhibits the pyruvate dehydrogenase in alcoholic subjects, so in our study pyruvate levels also decreased in ethanol subjects. Alcohol metabolism increases the NADH levels resulting to induce the lactate dehydrogenase activity (Phypher & Tom Pierce 2006). On the other hand alcohol metabolite acetaldehyde induced the insulin resistance to the cardiac tissue which reflect on variety metabolic abnormalities such as hyper lactacidemia, inhibit the lipolysis in adipocytes (Caballena, 2003) which resulting in decreased levels of the pyruvate in alcohol subjects. However with ginger treatment in ethanolic rats, carbohydrates and pyruvate levels are increased. Shanmugam *et al.*, (2009) have been established that ginger modulates the pyruvate and carbohydrate levels in diabetic rats. Our results were confined to above reports, due to the bioactive compounds in ginger, these compounds may increase the levels of carbohydrates and pyruvate in ethanol treated rats.

There was a significant increase ( $P < 0.05$ ) in the lactate and free amino acids levels in the ethanol treated rats. Conversely, the previous reports have been demonstrated that alcohol upshot on NADH/NAD<sup>+</sup> ratio which fallouts cellular redox state in turn increases lactate/ pyruvate ratio results in hyperlactacidemia (Greenway & Lutt 1990). Oxygen radicals and other ROS alter the proteins (Grune, 1997). These oxidative modifications may lead to changes in protein function, chemical fragmentation, or increased susceptibility to proteolytic attack (Freeman, 1982). The previous reports also demonstrated that chronic alcohol consumption can increases amino acids levels in alcoholic patients (Bleich *et al.*, 2004). Ginger intentionally condensed the proteolytic process and increases the gluconeogenesis (Chakraborty *et al.*,

2012), hence in ethanol treated rats with ginger treatment free amino acids levels are decreased.

Whereas ginger treatment decreases the levels of lactate and free amino acids compared to ethanol treatment group. Nevertheless, ginger has convalesced the cellular redox state by decreasing the NADH/NAD<sup>+</sup> ratio consequential reduced the free radicals by its antioxidant property (Shanmugam *et al.*, 2010). Dugasani *et al.*, (2010) have been reported that bioactive compounds of ginger like gingerol has antioxidant activity in various experimental modules. Furthermore, ginger extract has prevented free radical induced peroxidation of lipids, proteins and DNA (Morakinyo *et al.*, 2011). Similar consequences were may be occur in our study, so lactate and free amino acid levels are decreased in ginger treated group when compared to ethanol treated group.

### Conclusion

In the present study, the effect of ginger on carbohydrate metabolic profiles was established with reference to alcohol toxicity. Further research can be conducted by employing isolation of bioactive compounds and to test its role in alcohol induced myocardial infarction.

### Acknowledgement

This study was conducted with support of grants received from UGC-MRP (36-225/2008 (SR)).


### References



1. Ahmed, R., Sharma S (1997). Biochemical studies on combined effect of garlic (*Allium sativum* Linn) and ginger (*Zingiber officinale* Rosc) in albino rats. *Indian journal of experimental biology*, 35: 841-843.
2. American Psychiatric Association (1994).: *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed.; Washington, DC, USA.
3. Ajith TA, Nivitha V, Usha S (2007). *Zingiber officinale* Roscoe alone and in combination with alpha-tocopherol protect the kidney against cisplatin-induced acute renal failure. *Food Chem. Toxicol*, 45(6):921-927.
4. Baliunas, D.O., Taylor, B.J., Irving, H., Roerecke, M., Patra, J., Mohapatra, S., Rehm, J (2009) Alcohol as a risk factor for type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care*. 32: 2123-2132.
5. Banerjee B, D (1999) .The influence of various factors on immune toxicity assessment of pesticide chemicals. *Toxicol Lett*, 107: 21.
6. Bleich, S., Degner, D., Sperling, W., Bönsch, D., Thürauf, N., Kornhuber, J (2004). Homocysteine as a neurotoxin in chronic



- alcoholism. *Progress Neuropsychopharmacol. Biol. Psychiatr.*, 3: 453-464.
7. Bartsch, H., Nair, J (2000). Ultrasensitive and specific detection methods for exocytic DNA adducts; markers for lipid peroxidation and oxidative stress. *Toxicology*, 153: 105-114.
  8. Butt, M. S. and Sultan, M. T (2009). Green tea: Nature's Defense Against Malignancies. *Cri. Rev. Food Sci. Nutr.*, 49: 463-473.
  9. Caballena J. (2003). Current concepts in alcohol metabolism. *Ann Hepatol*, 2:60-66.
  10. Carroll N.V., Longley R.W. and Roe, J.H. (1956): Glycogen determination in liver and muscle by use of anthrone reagent. *J. Biol. Chem.*, 220:583-593.
  11. Cederbaum, A.I., Lu, Y., Wu, D. (2009). Role of oxidative stress in alcohol-induced liver injury. *Arch. Toxicol.*, 83:519-548.
  12. Chakraborty, D., Mukherjee, A., Sikdar, S., Paul, A., Ghosh, S., and Khuda-Bukhsa A. R., (2012.) "[6]-Gingerol isolated from ginger attenuates sodium arsenite induced oxidative stress and plays a corrective role in improving insulin signaling in mice," *Toxicology Letters*, 210(1): 34-43.
  13. Dugasani, S., Pichika, M. R.,Nadarajah,V. D., Balijepalli, M. K., Tandra, S.,Korlakunta, J. N. (2010).Comparative antioxidant and anti-inflammatory effects of [6]-gingerol, [8]-gingerol, [10]-gingerol and [6]-shogaol. *J. Ethnopharmacol.* 127: 515-520.
  14. Feinman, L., & Lieber, C.S (1998). Nutrition and diet in alcoholism. In: Shills, M.E., Oslon, J.A., Shike, M.and Ross, A.C; eds. *Modern Nutrition in Health and disease*. 9th ed. Baltimore: Williams and Wilkins. 1523-1542.
  15. Freeman BA., Crapo JD., (1982). Biology of disease: Free radicals and tissue injury. *Lab Invest.* 47:412-416.
  16. Friedmann J.E., and Hangen G.E. (1942): Pyruvic acid. Collection of blood for the determination of pyruvic and lactic acids. *J. Biol.Chem.*, 144: 67-77.
  17. George, A., Figueredo, V.M.,( 2010). Alcohol and arrhythmias: A comprehensive review. *J. Cardiov. Med. (Hagerstown)*, 11: 221-228.
  18. Greenway CV., Lautt WW (1990). Acute and chronic ethanol on hepatic oxygen ethanol and metabolism in cats. *Am J Physiol* 258:G411-G418.
  19. Grune T., Reinheckel T., Davies KJA (1997) . Degradation of oxidized proteins in mammalian cells. *FASEB J* 11:526-534.
  20. Guo, R., Ren, J (2010). Alcohol and acetaldehyde in public health: From marvel to menace. *Int. J. Environ. Res. Public Health*, 7: 1285-1301.
  21. Holford, N.H.G (1987).Clinical pharmacokinetics of ethanol. *Clin. Pharmacokinet.* 13:273-292.
  22. Huckabee WE. (1961) Relationship of pyruvate and lactate during anaerobic metabolism. V. Coronary adequacy. *Am J Physiol*, 200:1169-1176.
  23. Klein, H., Harmjan, D. (1975). Effect of ethanol infusion on the ultrastructure of human myocardium. *Postgrad. Med. J.*, 51: 325-329.
  24. Lieber CS. (1991). Hepatic metabolic and toxic effects of ethanol: 1991 update. *Alcohol Clin Exp Res*, 15:573-02.
  25. Lieber, CS, ed. (1995).Medical disorders of alcoholism. *N Engl J Med*,33 (16):1058-65.
  26. Lieber, C.S (2003). Relationships between Nutrition, Alcohol Use, and Liver disease. *Alcohol Health &Research World*,27 : 220-231.
  27. Lowry O.H., Rosebrough N.C., Farr A.L. and Randall R.J. (1951): Protein measurements with folin-phenol reagent. *Biol. Chem.*, 193: 265-275.
  28. Matias WG., Traore A., Bonini M., Sanni A.,Creppy EE.(1999).Oxygen reactive radicals production in cell culture by okadaic acid and their implication in protein synthesis inhibition. *Hum. Exp. Toxicol*, 18: 634-639.
  29. Morakinyo A.O., Akindele A J., Ahmed Z (2011).Modulation of Antioxidant Enzymes and Inflammatory Cytokines: Possible Mechanism of Anti-diabetic Effect of Ginger Extracts. *Afr. J. Biomed. Res.*, 14: 195 -202.
  30. Moore S., Stein WH (1954). Procedures for the chromatographic determination of amino acids on four per cent cross-linked sulfonated polystyrene resins. *J Biol Chem.* 211(2):893-906.
  31. Mukamal KJ., Jensen MK., Grønbaek M *et al* (2005). Drinking frequency, mediating biomarkers, and risk of myocardial infarction in women and men. *Circulation* 112:1406-1413.



32. Nicoll, R. and Henein, M. Y. (2009). Ginger (*Zingiber officinale Roscoe*): a hot remedy for cardiovascular disease? *Int. J. Cardiol.* **131**: 408–409.
33. Norberg, A., Jones, A.W., Hahn, R.G., Gabrielsson, J.L.(2003). Role of variability in explaining ethanol pharmacokinetics: Research and forensic applications. *Clin. Pharmacokinet*, **42**: 1-31.
34. Ohkubo, T., Metoki, H., Imai, Y. (2009). Alcohol intake, circadian blood pressure variation, and stroke. *Hypertension*, **53**: 4-5.
35. Phyper B., Tom Pierce JM (2006). Lactate physiology in health and disease. *The British Journal of Anaesthesia*, **6**: 3.
36. Randle P.J., Kerbey AL., Espinal J.(1988). Mechanisms decreasing glucose oxidation in diabetes and starvation: role of lipid fuels and hormones. *Diabetes Metab Rev* **4**: 623–638.
37. Regan, T.J (1990). Alcohol and the cardiovascular system. *JAMA*, **264**: 377-381.
38. Robin, M.A., Sauvage, I., Grandperret, T., Descatoire, V., Pessayre, D., Fromenty, B.(2005). Ethanol increases mitochondrial cytochrome P450 2E1 in mouse liver and rat hepatocytes. *FEBS Lett.* **579**: 6895-6902.
39. Rong, X., Peng, G., Suzuki, T., Yang, Q., Yamahara, J., and Li, Y. (2009). A 35-day gavage safety assessment of ginger in rats. *Regul. Toxicol. Pharmacol.* **54**:118–123.
40. Seitz, H.K., Becker, P. (2007). Alcohol metabolism and cancer risk. *Alcohol Res. Health*, **30**: 38-47.
41. Shanmugam, K.R., Ramkrishna, CH., Mallikarjuna, K., Sathyavelu Reddy K. (2010). Protective effect of ginger against alcohol induced renal damage and antioxidant enzymes in made albino rats, **48**:143-149.
42. Shanmugam, K.R., Ramakrishna, Ch., Mallikarjuna K., Sathyavelu Reddy, K (2009). The Impact of Ginger on Kidney Carbohydrate Metabolic Profiles in STZ Induced Diabetic Rats. *Asian J. Exp. Sc.*, **23** (1): 127-134
43. Siddaraju, M.N., Dharmesh, S.M. (2007). Inhibition of gastric H<sup>+</sup>-K<sup>+</sup> ATPase and Helicobacter pylori growth by phenolic antioxidants of *Zingiber officinale*. *Mol. Nutr. Food. Res.* **51**(3): 324-332.
44. Stadtman, ER (2004). Role of oxidant species in aging. *Current Medicinal Chemistry*, **11**: 1105–1112.
45. Van der Vusse GJ, Roemen THM, Prinzen F, Coumans WA., Reneman RS. (1982). Uptake and tissue content of fatty acids in dog myocardium under normoxic and ischemic conditions. *Circ Res*, **50**:538–546.
46. Vary TC., Lynch CJ., Lang CH (2001). Effect of chronic alcohol consumption on regulation of myocardial protein synthesis. *Am J. Physiol heart circ Physiol*, **281**:H242-51.
47. Young, HV, Luo, YL, Cheng, HY, Hsieh WC, Liao JC., Peng WC (2005). Analgesic and anti-inflammatory activities of [6]gingerol. *J Ethnopharmacol*, **96**: 207.
48. Zhang, Y., & Ren, J. (2010). Autophagy in ALDH2-elicited cardioprotection against ischemic heart disease: slayer or savior? *Autophagy*, **6**: 1212–1213.







Graph A-D: Changes in total carbohydrates (A) Pyruvate (B) Lactate (C) Proteins (D) Free amino acids (E) in cardiac tissue induced by ethanol administration and the effect of ginger treatment .Data mean  $\pm$  SD values (n=6) Groups that do not share same letters are significant ( $P < 0.05$ ) and letter with asterisk are more significant ( $P < 0.01$ ) to the control.