



Female ovarian hormone- Estriol's proconvulsant effect with kainic acid in Drosophila system model

Aakifa Ahmad*¹, Abdul Kalam Najmi² and Md. Naushad Alam³

1 Department of Pharmacology College of Pharmacy, Monad University, Hapur, Utter Pradesh, INDIA.

2 Department of Pharmacology, Faculty of Pharmacy, Jamia Hamdard, New Delhi, INDIA.

3 College of Pharmacy, IIIMT Greater Noida, Knowledge Park III, INDIA

Abstract

The effect of estriol would be evaluated for its effect on Drosophila fly model. Seizures in drosophila and man have several similarities and the utility of the fruit fly as a genetic model system for studying human seizures disorder and seizure susceptibility has clearly been demonstrated (Glasscock et al., 2005). Drosophila fly of 3-4 days old unmated fly were used. Drosophila model, optimized for screening of drugs dissolved in DMSO was used. Thirty flies were housed in each treatment vial. Three to four days old unmated male flies was treated with 8 mg/ml KA containing media for required time duration. A regimen for drosophila fly model where it was treated with kainic acid and resulted in increases as well as decrease of climbing speed for adult Drosophila fly. As higher organism has more colossal complexity that's why drosophila fly model is favored. This model is convenient to use and it is very relevant too.

Key-words: Estriol, seizures, drosophila fly model, climbing speed, kainic acid

Introduction

Seizures's association with estrogens are problematic to define due to complexity and heterogeneity of seizures. Contradictory data makes it more interesting subject regarding association of estrogen and epileptic seizures. General perception regarding estrogen's on seizures was till now that it increases neuronal excitability and mediates proconvulsant effects (Veliskiova,2006, Edwards et, al,1999, Nicoletti et al.1985) but estrogen has also exhibited anticonvulsant effect, this fact is supported by various animal and clinical data Veliskiova,2007, Reibel et al.,2000, Kalkbrenner and Standley, 2003, Tominaga et al.,2003). Estrogen is present in three biologically active form i.e estrone (E1), estradiol (E2) and estriol (E3). Most investigated form of estrogen is estradiol as it is present in post menopausal women.

Earlier studies of estradiol administration in rodents have revealed proconvulsant effects (Nicoletti et al., 1985). Estrogens applied to cortex could increase seizures (Marcus et al., 1966). In addition to this estradiol was shown to facilitate PTZ induced kindling (Lu et al., 2006), amygdala-kindled seizures (Buterbaugh, 1989) and kainic acid- induced seizures (Woolley, 2000) in adult female rats. Clinically also, it exacerbates seizures in women with epilepsy (Reddy, 2009).

Numerous similarities have been observed between seizures in drosophila fly and man (Glasscock et al.,2005, Hekmat-Scafe et al., Fergestad et al., 2006, Stilwell et al., 2006, Baraban, 2007, Song et al., 2007), this drosophila fly model is also useful in screening of anticonvulsant drug screening and drug target identification (Kuebler and Tanouye, 2002, Reynolds et al.,2004, Tan 2004, Hekmat-Scafe 2005). As higher organism has endless complexity, drosophila fly model is getting preference being simple organism (Joyce and Palsson, 2006). Inherent complexity of mammalian brain however does not render the available rodent epilepsy models as amenable to systems modeling (Gorter et al.,2006), because of the above reason we have selected drosophila fly model for understanding the mechanism underlying epileptogenesis. Seizures like activity in drosophila adults is known to be associated with loss of motor coordination and altered locomotor activities (Wang et al., 2004).

* Corresponding Author

Email: akfin.ahmad@gmail.com

Material and Methods

Animals

Drosophila melanogaster wild type Oregon-R strain was selected for the study. They were maintained at 24+1°C, 60% relative humidity and standard 12 hours light (9 AM to 9PM) and 12 hours dark cycle was maintained. The standard fly medium was used and it consisted of agar-agar, maize powder, dried yeast, nipagin and brown sugar. Standard methods of fly handling and manipulation were followed. Stringency required in behavioral studies was strictly adhered to at various levels, conditions of housing, light intensity. 3-4 days old unmated male flies were used and 30 flies were housed in each treatment vial. Institutional Animal Ethics Committee (IAEC) of Pinnacle Biomedical Research Institute (PBRI), Bhopal (Reg. No. 1824/PO/ERe/S/15/ CPCSEA) had approved the study and study approval reference number is PBRI/IAEC/PN-16026.

Drugs

Kainic acid, estriol and clomiphene citrate were purchased from Sigma, USA and diazepam was procured from Ranbaxy-Sun Pharma, India. Estriol was dissolved in dimethylsulfoxide (DMSO) and administered in doses of 0.005 and 0.01 mg/kg i.p. DMSO was given to control group as a vehicle. Clomiphene was dissolved in distilled water and administered intraperitoneally at dose of 0.9 mg/kg. Diazepam was taken as standard and given at 3 mg/kg.

EXPERIMENTAL DESIGN

Drosophila fly model methodology

Drosophila model, optimized for screening of drugs dissolved in DMSO, was used. Routine fly cultures was maintained at 24 ± 1°C, and 12 hrs light (9 AM to 9 PM) and 12 hrs dark cycle (Mohammad et al. 2009). Three to four days old unmated male flies was treated with 8 mg/ml KA containing media for required time duration. Thirty flies was housed in each treatment vial. After completion of 7 days, flies was placed in a 96-well plate and the movie was captured using a camera and the locomotor activity analyzed (Noldus Instruments). We took a glass column having 2 cm of internal diameter and 30 cm of length. Every fly was introduced for 90 seconds, dot/comma methods of measurement was applied, for purpose of monitoring locomotor activity of fly where the dot key depicts the moving fly and comma key depicts the resting fly. For calculation of climbing speed , the

formula is $s=h/t$ where s = climbing speed, h =height climmbed in cm, t = activity (seconds).

Table 1: Effect of various treatments on *Drosophila* fly model:

Following treatment groups was analyzed for locomotor behaviour change:

Groups	Treatment
1	Normal media containing 0.4% DMSO, 2 days, followed by normal media, 5 days
2	Normal media containing 8 mg/ml kainic acid and 0.4% DMSO, 2 days, followed by normal media containing 8 mg/ml kainic acid, 5 days
3	Normal media containing 8 mg/ml kainic acid and 0.4% DMSO in which the drug (Estriol, 5mM) was dissolved at 10 mM final concentration, 2 days, followed by normal media containing 8 mg/ml kainic acid, 5 days.
4	Normal media containing 8 mg/ml kainic acid and 0.4% DMSO in which the drug (Estriol, 10mM) was dissolved at 10 mM final concentration, 2 days, followed by normal media containing 8 mg/ml kainic acid, 5 days.
5	Normal media containing 8 mg/ml kainic acid and 0.4% DMSO, in which the drug (diazepam) was dissolved at 10 mM final concentration, 2 days, followed by normal media containing 8 mg/ml kainic acid, 5 days.

Results and Discussion

We observed through direct and indirect testing whether KA could produce effect in *drosophila* adults fly , although convulsions was not produced but flies treated with KA for 12 hours exhibited hyperactive movements. A decreased in "total directionality" was observed in KA treated flies compared to control (normal food). Flies treated with kainic acid took a serpent nous path, 16 mg/ml was the LC₅₀ dose (lethal data not presented). We substituted half the dose i.e 8 mg/ml for development and evaluating a chronic model.

We did not witnessed seizure like behavior in flies treated with 8 mg/ml, we followed a regimen by (Mohammad F et al., 2009) . They have developed a regimen where 7 days of chronic KA treatment followed by 7 days of absent KA treatment, it causes a decrease and increase in climbing speed in flies.

Due to immense complexity of higher organisms, approaches are currently focused on simpler organism (Joyce AR and Palsson , 2006, Chintapalli et al., 2007), thats why we included drosophila fly model in the study as the seizure in drosophila and man have several similarities and the utility of the fruit fly as a model system for studying human seizure disorders and seizure-susceptibility has clearly been established (Glasscock et al.,2005, Hekmat-Scafe et al., Fergestad et al., 2006, Stilwell et al., 2006, Baraban, 2007,Song et al., 2007).

Conclusion

As higher organism has more colossal complexity thats why drosophila fly model is favored. This model is convenient to use and it is very relevant too.

Acknowledgements

I am highly thankful to Pinnacle Biomedical Research Institute (Bhopal,India) for providing me with the opportunity to complete my Ph.D. research work.

References

1. Ahmad A and Vohora D. Proconvulsant effects of estriol, the third estrogen, in the mouse PTZ-kindling model. *Neuro Science*. 2014;10072-014-1795-4.
2. Baraban SC: Emerging epilepsy models: insights from mice, flies, worms and fish. *Curr Opin Neurol* 2007, 20:164-168. 23. Song J, Hu J, Tanouye M: Seizure suppression by top1 mutations in Drosophila. *J Neurosci* 2007, 27:2927-2937.
3. Barha, CK, Galea, LA. Influence of different estrogens on neuroplasticity and cognition in the hippocampus. *Biochimica Biophysica Acta*. 2010; 1800 : 1056- 67.
4. Brann DW, Dhandapani K, Wakade C, Virendra B, Khan MM. Neurotrophic and neuroprotective actions of estrogen: basic mechanisms and clinical implications. *Steroids*. 2007;72:381-405.
5. Brann, DW, Dhandapani, K, Wakade, C, Virendra, B, Mahesh, Khan, MM. Neurotrophic and neuroprotective actions of estrogen: Basic mechanisms and clinical implications. *Steroids*. 2007;72(5): 381-405.
6. Budziszewska B, Leśkiewicz M, Kubera M, Jaworska-Feil L, Kajta M, Lasoń W. Estrone, but not 17 beta-estradiol, attenuates kainate-induced seizures and toxicity in male mice. *Experimental Clinical Endocrinology and Diabetes*. 2001;109(3):168-73.
7. Charles V and Michael T. Morris water maze: procedures for assessing spatial and related forms of learning and memory. *Nature Protocol*. 2006; 2:848-858.
8. Check MD, Jerome H, Fred D, Lublin MD, Martin M, Mandel MD. Clomiphene as an Anticonvulsant Drug. *Archives of Neurology*. 1982;39(12):784.
9. Chintapalli VR, Wang J, Dow JAT: Using FlyAtlas to identify better Drosophila melanogaster models of human diseases. *Nat Genet* 2007, 39:715-720.
10. Draca S. Estriol and progesterone: a new role for sex hormones. *International Journal of Biomedical Science*. 2006; 2:305-307.
11. Edwards, HE, Bumham, WM, Mendonca, A, Bowlby, DA, MacLusky, NJ. Steroid hormones affect limbic after discharge thresholds and kindling rates in adult female rats. *Brain Research*. 1999;838: 2136-150.
12. Fergestad T, Bostwick B, Ganetzky B: Metabolic disruption in Drosophila bang-sensitive seizure mutants. *Genetics* 2006, 173:1357-1364.
13. Gibbs RB, Gabor R. Estrogen and cognition: applying preclinical findings to clinical perspectives. *Journal of Neuroscience Research*. 2003;74:637-643.
14. Glasscock E, Singhania A, Tanouye MA: The mei-P26 gene encodes an RBCC-NHL protein that regulates seizure susceptibility in Drosophila. *Genetics* 2005, 170:1677-1689.
15. Gorter JA, van Vliet EA, Aronica E, Breit T, Rauwerda H, Lopes da Silva FH, Wadman WJ: Potential new antiepileptic targets indicated by microarray analysis in a rat model for temporal lobe epilepsy. *J Neurosci* 2006, 26:11083-11110.
16. Hekmat-Scafe DS, Dang KN, Tanouye MA: Seizure suppression by gain-of-function escargot mutations. *Genetics* 2005, 169:1477-1493.
17. Hekmat-Scafe DS, Lundy MY, Ranga R, Tanouye MA: Mutations in the K⁺/Cl⁻cotransporter gene kazachoc (kcc) increase seizure susceptibility in Drosophila. *J Neurosci* 2006, 30:8943-8954.
18. Herzog AG. Seizure control with clomiphene therapy. A case report. *Archives of Neurology*. 1988; 45(2):209-10.


19. Joyce AR, Palsson BO: The model organism as a system: integrating 'omics' data sets. *Nat Rev Mol Cell Biol* 2006, 7:198-210.
20. Kalkbrenner, KA and Standley CA. Estrogen modulation of NMDA-induced seizures in ovariectomized and non-ovariectomized rats. *Brain Research*. 2003;28; 964: 244-9.
21. Kuebler D, Tanouye M: Anticonvulsant valproate reduces seizure-susceptibility in mutant *Drosophila*. *Brain Res* 2002, 958:36-42. 25.
22. Levin ER. Cellular Functions of the Plasma Membrane Estrogen Receptor. *Trends in Endocrinology and Metabolism*. 1999; 10(9): 374-377.
23. Marcus EM, Watson CW, Goldman PL. Effects of steroids on cerebral electrical activity: Epileptogenic effects of conjugated estrogens and related compounds in the cat and rabbit. *Archives Neurology*. 1966;15(5):521-532.
24. McEwen BS Invited Review: Estrogens effects on brain: Multiple sites and Molecular Mechanism. *Journal of Applied Physiology*. 2001; 91:2785-2801.
25. Mohammad F, Singh P, Sharma A. A *Drosophila* systems model of pentylenetetrazole induced locomotor plasticity responsive to antiepileptic drugs. 2009; 1752-0509-3-11.
26. Nicoletti F, Speciale C, Sortino MA, Summa G, Caruso G, Patti F, Canonico PL. Comparative effects of estradiol benzoate, the antiestrogen clomiphene citrate, and the progestin medroxyprogesterone acetate on kainic acid-induced seizures in male and female rats. *Epilepsia*. 1985; 26: 252-257.
27. Racine RJ. Modification of seizure activity by electrical stimulation II Motor seizure. *Electroencephalography Clinical Neurophysiology*. 1972; 32(3):281-94.
28. Reddy DS. The Role of Neurosteroids in the Pathophysiology and Treatment. *Epilepsy Research*. 2009;85(1): 1-30.
29. Reibel S, Vivien-Roels B, Le BT, Larmet Y, Carnahan J, Marescaux C, Depaulis A. Overexpression of neuropeptide Y induced by brain-derived neurotrophic factor in the rat hippocampus is long lasting. *European Journal Neuroscience*. 2000;12(2):595-605.
30. Reynolds ER, Stauffer EA, Feeney L, Rojahn E, Jacobs B, McKeener C: Treatments with the antiepileptic drugs phenytoin and gabapentin ameliorates seizure and paralysis of *Drosophila* bang-sensitive mutants. *J Neurobiol* 2004, 58:503-513.
31. Scharfman HE, Mercurio TC, Goodman JH, Wilson MA, MazLusky NJ. Hippocampal excitability increases during the estrous cycle in the rat: a potential role for brain-derived neurotrophic factor. *Journal Neuroscience*. 2003;23:11641-11652.
32. Stilwell GE, Saraswati S, Littleton JT, Chouinard SW: Development of a *Drosophila* seizure model for *in vivo* high-throughput drug screening. *Eur J Neurosci* 2006, 24:2211-2222.
33. Takahashi K, Okada M, Ozaki T, Kurioka H, Manabe A, Kanasaki H, Miyazaki K. Safety and efficacy of estriol for symptoms natural or surgically induced menopause. *Human Reproduction*. 2000;15:1028-1036.
34. Tan JS, Lin F, Tanouye MA: Potassium bromide, an anticonvulsant, is effective at alleviating seizures in the *Drosophila* bang-sensitive mutant bang-senseless. *Brain Res* 2004, 1020:45-52. 27.
35. Tan RS, Pu SJ, Culberson JW. Role of androgens in mild cognitive impairment and possible interventions during andropause. *Medical Hypotheses*. 2003; 60:448-452.
36. Tominaga K, Yamauchi A, Shuto H, Niizeki M, Makino K, Oishi R, Kataoka Y. Ovariectomy aggravates convulsions and hippocampal gamma-aminobutyric acid inhibition induced by cyclosporin A in rats. *European Journal Pharmacology*. 2003; 24: 5,675-704.
37. Veliskova J. The role of estrogen in seizures and epilepsy: the bad guys or the good guys? *Neuroscience*. 2006;138: 837-844.
38. Veliskova J. Estrogens and Epilepsy: why are we so excited? *Neuroscientist*. 2007; 13:77-88.
39. Vohora D, Pal SN, Pillai KK. Modulation of spontaneous alternation behavior of mice treated with thioperamide and tacrine in a cross maze task. *Fundamental & Clinical Pharmacology*. 2005;19(5):531-2.

40. Wahnschaffe U, Löscher W.) Lack of changes in seizure susceptibility during the estrous cycle in kindled rats. *Epilepsy Research.* 1992;13(3):199-204.

41. Wang P, Saraswati S, Guan Z, Watkins CJ, Wurtman RJ, Littleton JT: A Drosophila Temperature-Sensitive Seizure Mutant in Phosphoglycerate Kinase disrupts ATP Generation and Alters Synaptic Function. *J Neurosci* 2004, 24:4518-4529.


42. Zemlyak I, Sheila MB, Robert MS. Protection against gp120-induced neurotoxicity by an array of estrogenic steroids. *Brain Research.* 2002; 958:272-276.

Figure 1: Treatment with chronic KA and causes changes in climbing speed when flies are treated with normal food.




All the values were expressed as mean \pm SEM. A treatment with kainic acid for flies treated with normal food.

Figure 2: Absence of KA causes changes in climbing speed when flies are treated with normal food




All the values were expressed as mean \pm SEM.

Figure 3: Climbing speed was recorded on 12 hours, 2nd day and 7th day after KA treatment and then on 3rd day, 5th day and 7th day after KA with drawl.

All the values were expressed as mean \pm SEM. Climbing speed was recorded on 12 hours, 2nd day and 7th day after KA treatment and then on 3rd day, 5th day and 7th day after KA with drawl. For control climbing assay to examine consistency among normal food flies housed in different vials. For stats we applied ANOVA. * indicates significant difference between control (normal food) and KA treatment. *p* values are provided over asterisks.

Figure 4: Effect of various treatments on drosophila fly model on respective time.



All the values were expressed as mean \pm SEM. VEH: Vehicle (Dimethyl Sulfoxide); EST: Estriol; CLO: Clomiphene; DZP: Diazepam; KA: Kainic acid; KA (25 mg/kg) was administered once every 2 days for 5 weeks while EST 1 (0.005 mg/kg) and EST 2 (0.01 mg/ kg) and CLO (0.9 mg/kg) were administered daily.

How to cite this article

Ahmad A., Najmi A.K. and Alam Md. N. (2019). Female ovarian hormone- Estriol's proconvulsant effect with kainic acid in Drosophila system mode. *Int. J. Pharm. Life Sci.*, 10(7-8):6342-6348.

Source of Support: Nil; Conflict of Interest: None declared

Received: 27.05.19; Revised: 20.06.19; Accepted: 22.07.19